

2019

## The relationship between body mass index (BMI) and sedentary behavior is mediated by negative peer interaction in boys

Jacob E. Barkley  
*Kent State University*

Gregory S. Farnell  
*John Carroll University*

Follow this and additional works at: [https://collected.jcu.edu/fac\\_bib\\_2019](https://collected.jcu.edu/fac_bib_2019)




Part of the [Health and Physical Education Commons](#)

---

### Recommended Citation

Barkley, Jacob E. and Farnell, Gregory S., "The relationship between body mass index (BMI) and sedentary behavior is mediated by negative peer interaction in boys" (2019). *2019 Faculty Bibliography*. 14.  
[https://collected.jcu.edu/fac\\_bib\\_2019/14](https://collected.jcu.edu/fac_bib_2019/14)

This Article is brought to you for free and open access by the Faculty Bibliographies Community Homepage at Carroll Collected. It has been accepted for inclusion in 2019 Faculty Bibliography by an authorized administrator of Carroll Collected. For more information, please contact [mchercourt@jcu.edu](mailto:mchercourt@jcu.edu).



## The Relationship between Body Mass Index (BMI) and Sedentary Behavior is Mediated by Negative Peer Interaction in Boys.

Received 01<sup>st</sup> January 2019  
Accepted 12<sup>th</sup> March 2019

[www.ijpefs.com](http://www.ijpefs.com)

Jacob E. Barkley<sup>a,\*</sup>, Gregory S. Farnell<sup>b</sup>,

<sup>a</sup> Associate Professor, School of Health Sciences, Kent State University, Ohio.

<sup>b</sup> Associate Professor, John Carroll University, Department of Exercise Science & Sports Studies, Ohio.

\* Corresponding Author: Ph: 330-672-0209; Email: [jbarkle1@kent.edu](mailto:jbarkle1@kent.edu)

**Abstract:** To determine if self-reported negative social interaction mediates the relationship between sedentary behavior and body mass index (BMI) percentile in boys. Twelve overweight/obese ( $\geq 85^{\text{th}}$  BMI percentile) and 14 non-overweight ( $< 85^{\text{th}}$  BMI percentile) boys ( $10.5 \pm 1.5$  years old) completed surveys assessing overt peer victimization and relational victimization. Children were individually given access to a gymnasium with physical activity equipment and sedentary alternatives for 30 minutes. Children could play with the equipment in any pattern they wished and the amount of time allocated to sedentary activities (sitting time) was recorded. Overt and relational victimization were moderately and positively associated with BMI percentile ( $r \geq 0.40$ ,  $p \leq 0.04$ ) and sitting time ( $r \geq 0.40$ ,  $p \leq 0.05$ ) and sitting time was positively associated with BMI percentile ( $r = 0.4$ ,  $p = 0.05$ ). After controlling for overt and relational victimization the correlation between sitting time and BMI percentile was non-significant ( $r \leq 0.28$ ,  $p \geq 0.18$ ). The positive relationship between BMI percentile and sedentary behavior was mediated by measures of negative social interaction.

**Key Words:** Child, peer influence, peer victimization.

### 1. Introduction

Overweight/obese children allocate more time to sedentary behaviour (i.e., sitting) and are less physically active than their non-overweight peers [1-6]. This phenomenon has been demonstrated in free-living environments using self-report surveys as well as objective physical activity monitors (e.g., accelerometers) [7-9]. There is also evidence of disparate sedentary behaviour and physical activity between overweight/obese and non-overweight children in controlled laboratory environments using objective measures. [10,11] Several factors have been identified that may explain this disparity including differences between overweight/obese and non-overweight children in: the relative reinforcing (i.e., motivating) value of physical activity versus sedentary alternative, self-efficacy for physical activity, self-confidence and discomfort during

exercise [12-18].

In addition to the variables outlined above, negative social interaction has been identified as a potentially important factor in explaining why overweight/obese children are less active and more sedentary than their non-overweight peers [4, 7, 11, 19-23]. Overweight/obese children are more frequently the target of negative social interaction than their non-overweight peers [4, 10, 11, 24, 25]. This negative social interaction can include less social support and greater overt and relational victimization [26-29]. Overt victimization are threats or acts of physical violence. Relational victimization is attempts by peers to harm a child's relationships with other children. These forms of negative social interaction are associated with greater sedentary behaviour and reduced physical activity in children [4, 7, 11]. Additionally, ostracism (i.e., social

exclusion) is also more common among overweight/obese children and there is experimental evidence suggesting that a bout of simulated ostracism causes a subsequent reduction in physical activity and increase in sedentary behaviour in children regardless of their bodyweight.<sup>30</sup> Taken together these findings suggest that overweight/obese children are more prone to negative social interaction and this negative social interaction is predictive of reduced physical activity and greater sedentary behaviour. This suggests that negative social interaction is a potential mediator of the relationship between adiposity and physical activity/sedentary behaviour. However, the ability of negative social interaction to mediate the relationship between adiposity and physical activity/sedentary behaviour in youth has not been previously tested.

Therefore, the purpose of this study was to assess the relationships between self-reported measures of peer victimization (overt and relational victimization), body mass index (BMI) percentile for age and objectively-observed sedentary behavior during free play in a controlled laboratory setting in boys. We then assessed the ability of overt and relational victimization to mediate the relationship between BMI percentile and sedentary behavior. We hypothesized that both measures of peer victimization, BMI percentile and sedentary behavior would all be significantly and positively related to one another. In other words, children with a greater BMI percentile would report greater peer victimization and participate in greater sedentary behavior than non-overweight children and greater peer victimization would be associated with greater sedentary behavior. We also hypothesized that the significant, positive relationship between BMI percentile and sedentary behavior would be rendered non-significant when controlling for peer victimization, thus identifying peer victimization as a potential mediator of this relationship.

## 2. Methods

Participants included 14 non-overweight (BMI <85<sup>th</sup> percentile) boys (10.1 ± 1.4 years old, 138.7 ± 9.5 cm, 33.5 ± 6.2 kg, 56.2 ± 15.8 BMI

percentile) and 12 overweight/obese (BMI >85<sup>th</sup> percentile) boys (10.8 ± 1.6 years old, 147.6 ± 9.6 cm, 56.8 ± 17.1 kg, 93.4 ± 6.0 BMI percentile). Participants were recruited from a database of families who had previously participated in unrelated studies in our laboratory and from flyers posted in the local community. All participants were free from any orthopedic, cardiovascular, metabolic or cognitive disorders that would prevent them from safely participating in physical activity. Participants and a parent/legal guardian read and signed assent and consent forms, respectively. All procedures were approved by the University institutional review board.

### 2.1 Procedure

Children completed a single laboratory/activity session. While in the laboratory, they were measured for height and weight using a digital stadiometer (Charder, Taichung City, Taiwan) and balance beam scale (Health O Meter, Alsip, IL), respectively. Children also completed the validated Children Self-Experience Questionnaire to assess self-reported incidence of peer victimization [26-27]. Children then participated in physical activity and sedentary behavior in any pattern they chose during a 30-minute, free-play session in a 4,300 square foot gymnasium that was located within the same facility as the laboratory. The gymnasium was equipped with a variety of physical activities (obstacle courses, balls, hoops, etc.) and a table equipped with a chair and sedentary activities (age-appropriate books, toys, coloring sheets, crayons, pencils, etc.). The specific configuration of the gymnasium and activity options has been reported previously [31-32]. Each child participated in this free-play activity session with no other children present as the presence of a peer can affect physical activity and sedentary behavior [4, 10, 32]. During this session, research personnel recorded the time children allocated to the sedentary activities. If children wished to play with the sedentary activities, they were instructed they had to do so while seated in the chair located at the table with said activities.

### 2.2 Measurements

**Peer victimization:** Children completed the

validated Children Self-Experience Questionnaire to assess peer victimization. This questionnaire reports peer victimization in two different subscales: overt and relational victimization. The overt victimization subscale assessed the frequency of threats or acts of physical violence a child is subjected to by their peers. The relational victimization subscale assessed the frequency of attempts to harm relationships children were subjected to by their peers. These scales have previously been shown to possess strong internal consistency (Cronbach's  $\alpha = 0.82 - 0.97$ ) [26, 27, 29, 33].

**Sedentary behavior:** The amount of time children allocated to sedentary behavior during the 30-minute activity session was recorded via a stopwatch (Traceable® Stopwatch, Fisher Scientific, Waltham, Massachusetts) by research personnel discretely observing the participant. The stopwatch was started at the moment a child sat in the chair at the sedentary activity table and stopped when/if they vacated the chair and returned to the physical activities. The process was repeated should a child participate in multiple bouts of sedentary activity during the 30-minute activity session. The time allocated to these multiple bouts of sedentary behavior was then summed as the measure of sedentary behavior.

### 2.3 Analytic plan

Independent samples t-tests were utilized to assess potential differences in BMI percentile, self-reported peer victimization (overt, relational) and sedentary behavior in non-overweight and overweight boys. The remaining analytic approach was designed to assess the ability of the two scales of self-reported peer victimization to mediate the relationship between BMI percentile and sedentary behavior. According to Baron and Kenny, a mediator

is a variable that accounts for the relationship between two other variables [34]. This mediator will be correlated to both of the other two variables and the correlation between these two other variables will be rendered non-significant after controlling for the potential mediator. Therefore Pearson's correlation analyses were first performed to assess the relationship between the following variables: peer victimization subscales (overt, relational), BMI percentile and sedentary time. Subsequent partial correlations were then performed assessing the relationship between BMI percentile and sedentary time after individually controlling for the two peer victimization subscales

### 3. Results


Mean comparisons between non-overweight and overweight boys are listed in Table 1.

Peer victimization subscales (overt, relational) were significantly and positively correlated ( $r = 0.85, p < 0.001$ ) to one another and each subscale was significantly and positively correlated to both BMI percentile ( $r = 0.46, p = 0.02$  for overt,  $r = 0.40, p = 0.04$  for relational) and sedentary behavior ( $r = 0.40, p = 0.05$  for overt,  $r = 0.42, p = 0.04$  for relational). In other words, children reporting greater peer victimization had a greater BMI percentile and participated in more sedentary behavior. BMI percentile was also significantly and positively correlated to sedentary behavior ( $r = 0.4, p = 0.05$ ). In other words, children with a greater BMI percentile participated in more sedentary behavior.

Partial correlations revealed that when separately controlling for each of the two peer victimization subscales the relationship between BMI percentile and sedentary behavior was rendered non-significant ( $r = 0.28, p = 0.18$  for overt,  $r = 0.26, p = 0.21$  for relational, Figure 1).

**Table 1.** Data are means  $\pm$  SD. There were significant differences ( $t \geq 2.3, p \leq 0.03$ ) between groups for all variables.

	BMI percentile	Overt victimization	Relational victimization	Sedentary time (min)
Non-overweight boys	56.2 $\pm$ 15.8	7.7 $\pm$ 2.6	8.2 $\pm$ 3.2	1.4 $\pm$ 3.6
Overweight/obese boys	93.4 $\pm$ 6.0	11.5 $\pm$ 3.9	11.8 $\pm$ 4.7	7.7 $\pm$ 6.6



**Figure 1.** Illustrates the correlations before (A) and after separately controlling for the relational victimization (B) and overt victimization (C) subscales of the peer victimization questionnaire.

#### 4. Discussion

The purpose of this study was to assess whether or not self-reported measures of peer victimization mediated the relationship between BMI percentile and sedentary behavior in boys during free play in a controlled environment. Presently, overweight boys reported greater peer victimization and were more sedentary during the 30-minute activity session than non-overweight boys. There were also positive relationships between BMI percentile, sedentary behavior and both scales of self-reported peer victimization. However, when separately controlling for each of the two peer victimization scales, the correlations between BMI percentile and sedentary behavior were no longer significant. According to the methodology proposed by Baron and Kenny, self-reported peer victimization did mediate the relationship between BMI percentile and sedentary behavior [34]. In other words, peer victimization may play an important role in predicting the greater sedentary behavior seen in overweight/obese youth.

Previous research from our group and others have reported that children who experience more negative social interaction (e.g., peer victimization) are less physically active and more sedentary in both controlled and free-living environments than peers who experience less negative interaction [4,7,11,19-23]. We have also demonstrated that a bout of negative peer interaction (i.e., simulated ostracism) causes a subsequent reduction in physical activity and an increase in sedentary behavior. Furthermore, we and others have reported that overweight/obese children are more likely to be the targets of negative peer interaction and are less physically active than their non-overweight peers [4,10,11,24-25]. Therefore, it is possible that the more frequent negative peer interaction reported by overweight/obese youth may be, at least in part, an explanatory factor behind why these overweight/obese children are more sedentary and less physically active than their non-overweight peers. The present finding that peer victimization mediated the relationship between BMI percentile and sedentary behavior in boys supports this notion.

The ability of peer victimization to mediate the relationship between BMI percentile and sedentary behavior in boys is a novel finding. However, research from Storch et al offers insight into the possible mechanism behind the effect negative peer interaction may have on physical activity/sedentary behavior in children.<sup>19</sup> Similar to the present study, Storch et al used Baron and Kenny's approach and reported that symptoms of depression mediated the negative relationship between peer victimization and physical activity. It is therefore possible that depression, possibly as a result of peer victimization, suppresses physical activity. Taking this and the present findings together, there are relationships between being overweight/obese, increased peer victimization, greater depressive symptoms and reduced physical activity/increased sedentary behavior. If this is true, suppressed physical activity/greater sedentary behavior may then contribute to greater depression and weight gain which would likely lead to further negative peer interaction [24-25, 35-38] In other words, the relationship between elevated BMI percentile, negative peer interaction, greater depression and physical inactivity/greater sedentary behavior may be cyclical.

While this study provides additional evidence of the role that negative peer interaction may play in children's sedentary behavior, it is not without limitations. First, the study examines a small sample and only included boys. Future research should include a larger sample and also examine girls. However, while there is evidence that certain types of negative peer interaction (e.g., weight criticism) may be more prevalent in girls than boys, prior research examining the impact of peer victimization on physical activity and sedentary behavior in both boys and girls has found no differences between the sexes.<sup>7,19,30</sup> Second, while the purpose of the study was to examine sedentary behavior in a controlled, free-play environment, future research should include separate measures of physical activity (e.g., accelerometry), apart from time allocated to activities. Because physical activity and sedentary behavior, while typically correlated to one another, are independent risk factors of a myriad of cardio-metabolic disorders, it is worthwhile to assess both

variables [6, 39-41]. Finally, measures of peer victimization were self-reported thus removing the ability to make causal inferences regarding the effect of peer victimization upon sedentary behavior and BMI percentile. However, prior experimental research has reported that simulated negative peer interaction caused a subsequent increase in children's sedentary behavior regardless of the child's sex or BMI percentile [30].

## 5. Conclusion

In conclusion, presently self-reported peer victimization mediated the positive relationship between BMI percentile and sedentary behavior in boys during free play in a controlled environment. This finding is in support of previous experimental and non-experimental research indicating negative peer interaction may cause and/or is associated with increased sedentary behavior and reduced physical activity. Taken together, there is mounting evidence of the importance of positive peer interaction to promote physical activity and discourage sedentary behavior in children. Additional research examining this mediating effect of peer victimization in girls is warranted.

## References

- [1] S.J. te Velde, I. De Bourdeaudhuij, I. Thorsdottir, Mette Rasmussen, Maria Hagströmer, Knut-Inge Klepp, Johannes Brug, Patterns in sedentary and exercise behaviors and associations with overweight in 9–14-year-old boys and girls - a cross-sectional study, *BMC Public Health*, 7 (2007) 16.
- [2] J. Crespo, E. Smit, R.P. Troiano, S.J. Bartlett, C.A. Macera, R.E. Andersen Television watching, energy intake, and obesity in US children: results from the third National Health and Nutrition Examination Survey, 1988-1994, *Archives of Pediatrics and Adolescent Medicine*, 155 (2001) 360-365.
- [3] K.S. Steinbeck, The importance of physical activity in the prevention of overweight and obesity in childhood: a review and an opinion, *Obesity Reviews*, 2(2001) 117-130.

- [4] S.J. Salvy, J.C. Bowker, L. Germeroth, J. Barkley, Influence of Peers and Friends on Overweight/Obese Youths' Physical Activity, *Exercise and Sport Sciences Reviews*, 40 (2012) 127-132.
- [5] V. Carson, S. Hunter, N. Kuzik, C.E. Gray, V.J. Poitras, J.P. Chaput, T.J. Saunders, P.T. Katzmarzyk, A.D. Okely, S. Connor Gorber, M.E. Kho, M. Sampson, H. Lee, Tremblay M.S. Systematic review of sedentary behaviour and health indicators in school-aged children and youth: an update, *Applied Physiology Nutrition and Metabolism*, 41 (2016) 240-265.
- [6] A. Must, D.J. Tybor Physical activity and sedentary behavior: a review of longitudinal studies of weight and adiposity in youth, *International journal of obesity and related metabolic disorders*, 29 (2005) 84-96.
- [7] M.S. Faith, M.A. Leone, T.S. Ayers, M. Heo, A. Pietrobelli, Weight criticism during physical activity, coping skills, and reported physical activity in children, *Pediatrics*, 110 (2002) 23.
- [8] V. Carson, M.S. Tremblay, J.P. Chaput, S.F. Chastin, Associations between sleep duration, sedentary time, physical activity, and health indicators among Canadian children and youth using compositional analyses, *Applied Physiology Nutrition and Metabolism*, 41 (2016) 294-302.
- [9] V. Cleland, D. Crawford, L.A. Baur, C. Hume, A. Timperio, J. Salmon, A prospective examination of children's time spent outdoors, objectively measured physical activity and overweight, *International Journal of Obesity*, 32 (2008) 1685-1693.
- [10] M. Rittenhouse, S.J. Salvy, J.E. Barkley, The effect of peer influence on the amount of physical activity performed in 8- to 12-year-old boys, *Pediatric Exercise Science*, 23 (2011) 49-60.
- [11] M. Rittenhouse, J. Barkley, Self-reported peer victimization and objectively measured physical activity behaviour in boys: a quasi-experimental study, *Journal of Exercise Physiology*, 16 (2013) 84-92.
- [12] J.F. Sallis, J.J. Prochaska, W.C. Taylor, J.O. Hill, J.C. Geraci, Correlates of physical activity in a national sample of girls and boys in grades 4 through 12, *Health Psychology*, 18 (1999) 410-415.
- [13] J.F. Sallis, J.J. Prochaska, W.C. Taylor, A review of correlates of physical activity of children and adolescents, *Medicine & Science in Sports & Exercise*, 32 (2000) 963-975.
- [14] L.H. Epstein, R.A. Paluch, K.J. Coleman, D. Vito, K. Anderson, Determinants of physical activity in obese children assessed by accelerometer and self-report, *Medicine & Science in Sports & Exercise*, 28 (1996) 1157-1164.
- [15] L.H. Epstein, J.A. Smith, L.S. Vara, J.S. Rodefer, Behavioral economic analysis of activity choice in obese children, *Health Psychology*, 10 (1991) 311-316.
- [16] S.G. Trost, L.M. Kerr, D.S. Ward, R.R. Pate, Physical activity and determinants of physical activity in obese and non-obese children, *International journal of obesity and related metabolic disorders*, 25 (2001) 822-829.
- [17] J.W. Pierce, J. Wardle Cause, effect beliefs and self-esteem of overweight children, *Journal of Child Psychology and Psychiatry*, 38 (1997) 645-650.
- [18] S.M. Smith, B. Sumar, K.A. Dixon, Musculoskeletal pain in overweight and obese children, *International Journal of Obesity*, 38 (2014) 11-15.
- [19] E.A. Storch, V.A. Milsom, N. Debraganza, A.B. Lewin, G.R. Geffken, J.H. Silverstein, Peer victimization, psychosocial adjustment, and physical activity in overweight and at-risk-for-overweight youth, *Journal of Pediatric Psychology*, 32 (2007) 80-89.
- [20] M. Hohepa, R. Scragg, G. Schofield, G.S. Kolt, D. Schaaf, Social support for youth physical activity: Importance of siblings, parents, friends and school support across a segmented school day, *International Journal of Behavioural Nutrition and Physical Activity*, 4 (2007) 54.
- [21] M.A. Kunesch, C.A. Hasbrook, R. Lewthwaite, Physical activity socialization: Peer interactions and affective responses among a sample of sixth grade girls, *Sociology of Sport Journal*, 9 (1992) 385-396.
- [22] R.M. Page, J. Frey, R. Talbert, C. Falk, Children's

- feelings of loneliness and social dissatisfaction: Relationship to measures of physical fitness and activity, *Journal of Teaching in Physical Education*, 11(1992) 211-219.
- [23] A.L. Smith, Perceptions of peer relationships and physical activity participation in early adolescence, *Journal of Sport and Exercise Psychology*, 21(1999) 329-350.
- [24] D. Neumark-Sztainer, N. Falkner, M. Story, C. Perry, P.J. Hannan, S. Mulert, Weight-teasing among adolescents Correlations with weight status and disordered eating behaviors, *International journal of obesity and related metabolic disorders*, 26 (2002) 123-131.
- [25] H.A. Hayden-Wade, R.I. Stein, A. Ghaderi, B.E. Saelens, M.F. Zabinski, D.E. Wilfley, Prevalence, characteristics, and correlates of teasing experiences among overweight children vs. non-overweight peers, *Obesity Research*, 13 (2005) 1381-1392.
- [26] N.R. Crick, J.K. Grotpeter, Children's treatment by peers: Victims of relational and overt aggression, *Development and Psychopathology*, 8 (1996) 367-380.
- [27] J.K. Grotpeter, N.R. Crick, Relational Aggression, Overt Aggression, and Friendship, *Child development*, 67 (1996) 2328-2338.
- [28] K.D. Williams, Ostracism: The Kiss of Social Death, *Social and Personality Psychology Compass*, 1 (2007) 236-247.
- [29] N.R. Crick, M.A. Bigbee, Relational and overt forms of peer victimization: a multiinformant approach, *Journal of Consulting and Clinical Psychology*, 66 (1998) 337-347.
- [30] J.E. Barkley, S.J. Salvy, J.N. Roemmich, The effect of simulated ostracism on physical activity behaviour in children, *Pediatrics*, 129 (2012) 659-666.
- [31] G. Sanders, C. Peacock, M.L. Williamson, K. Wilson, A. Carnes, J.E. Barkley, The effect of friendship groups on children's physical activity: An experimental study, *Journal of Behavioural Health*, 3 (2014) 95-100.
- [32] J.E. Barkley, S.J. Salvy, G.J. Sanders, S. Dey, K.P. Von Carlowitz, M.L. Williamson, Peer Influence and Physical Activity Behaviour in Young Children: An Experimental Study, *Journal of Physical Activity and Health*, 11 (2013) 404-409.
- [33] N.R. Crick, J.F. Casas, M. Mosher Relational and overt aggression in preschool, *Developmental psychology*, 33(1997) 579-588.
- [34] R.M. Baron, D.A. Kenny, The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations, *Journal of Personality and Social Psychology*, 51 (1986) 1173-1182.
- [35] P. Kremer, C. Elshaug, E. Leslie, J.W. Toumbourou, G.C. Patton, J. Williams, Physical activity, leisure-time screen use and depression among children and young adolescents, *Journal of Science and Medicine in Sport*, 17 (2014) 183-187.
- [36] G.M. Reeves, T.T. Postolache, S. Snitker, Childhood Obesity and Depression: Connection between these Growing Problems in Growing Children, *International Journal of Child Health and Human Development*, 1 (2008) 103-114.
- [37] S.J. Salvy, J.W. Bowker, J.N. Roemmich, R.D. Natalie Romero, M.A. Elizabeth Kieffer, M.S. Rocco Paluch, L. H. Epstein, Peer Influence on Children's Physical Activity: An Experience Sampling Study, *Journal of Pediatric Psychology*, 33 (2008) 39-49.
- [38] K.M.J. Lagerspetz, K. A.J. Björkqvist, M. Berts, E. King, Group aggression among school children in three schools, *Scandinavian Journal of Psychology*, 23 (1982) 45-52.
- [39] S.J. Marshall, S.J.H. Biddle, T. Gorely, N. Cameron, I. Murdey Relationships between media use, body fatness and physical activity in children and youth: a meta-analysis, *International journal of obesity and related metabolic disorders*, 28 (2004) 1238-1246.
- [40] M.F. Hjorth, J.P. Chaput, C. Ritz, S.M. Dalskov, R. Andersen, A. Astrup, I. Tetens, K.F. Michaelsen, A. Sjödin, Fatness predicts decreased physical activity and increased sedentary time, but not vice versa: support from a longitudinal study in 8- to 11-year-old children, *International Journal of Obesity*, 38 (2014) 959-965.
- [41] U. Ekelund, S. Brage, K. Froberg, M. Harro, S.A.



Anderssen, L.B. Sardinha, C. Riddoch, L.B. Andersen, TV Viewing and Physical Activity Are Independently Associated with Metabolic Risk in Children: The European Youth Heart Study, *PLOS Medicine*, 3 (2006) 488.

### Acknowledgements

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors

**Competing Interests:** The author declares to have no competing interests

### About The License



The text of this article is licensed under a Creative Commons Attribution 4.0 International License